

PEGO

BIOSTIMULANT OF FLOWERING AND FRUIT DEVELOPMENT ORGANIC NITROGEN FERTILIZER Extract of yeast with brown seaweed, calcium, boron and zinc

COMPOSITION (%w/w)

| Organic nitrogen (N) | 1% |
|--|-------|
| Water-soluble calcium oxide (CaO) | 5% |
| Water-soluble boron (B) | 0.2% |
| Water-soluble zinc (Zn) | 0.15% |
| Zinc (Zn) chelated with EDTA | 0.15% |
| Organic biological carbon (C) | 10% |
| рНб | |
| Organic matter with molecular weight <50 kDa | 30% |
| | |

PEGO is a formulation of both organic and mineral nature. Its special composition makes it ideal for promoting and catalyzing the fruit development. The organic components are particularly effective because they contain specific elements such as propolis, which allow to attract a greater number of pollinator (bees and bombs) and essential amino acids that favor the absorption of the calcium, boron and zinc components, which are vital for flowering and fruit attachment. The application of PEGO, therefore enables flower pollination, fruits attachment and it results in abundant harvests of excellent quality.

| DOSAGE AND METHODS OF USE | CROPS | APPLICATION | DOSE |
|------------------------------|---|------------------------------------|------------------------------|
| FOLIAR | Fruit Horticultural Floricultural Industrial | 2-3 Before and during flowering | 200-250 ml/100 L of water |
| FERTIGATION | Fruit Horticultural Floricultural Industrial | 2-3 Before and during flowering | 5-8 L/ha |

Warnings: Shake the product well before use. For any mixtures with other products, make preliminary tests. Observe the dosages shown. The product is not combustible and is stable at ordinary temperatures and pressures. Store the product in adequately ventilated rooms at temperatures between +5 ° C and +35 ° C.

| FORMULATION | PACKAGE |
|----------------|---------|
| Soluble liquid | |